mH2A1 Conjugated Antibody

Catalog No: #C30801



 Package Size:
 #C30801-AF350 100ul
 #C30801-AF405 100ul
 #C30801-AF488 100ul

 #C30801-AF555 100ul
 #C30801-AF594 100ul
 #C30801-AF647 100ul

 #C30801-AF680 100ul
 #C30801-AF750 100ul
 #C30801-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	mH2A1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	lgG
Purification	Affinity purification
Applications	most applications
Species Reactivity	Hu,Ms,Rt
Immunogen Description	Recombinant fusion protein of human H2AFY (NP_613258.2).
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	H2AFY; H2A.y; H2A/y; H2AF12M; MACROH2A1.1; mH2A1; macroH2A1.2; core histone macro-H2A.1
Accession No.	Swiss-Prot#:075367NCBI Gene ID:9555
Uniprot	O75367
GeneID	9555;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	40kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:
AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. It replaces conventional H2A histones in a subset of nucleosomes where it represses transcription and participates in stable X chromosome inactivation. Alternative splicing results in multiple transcript variants encoding different isoforms.

Note: This product is for in vitro research use only