

# WFDC3 Conjugated Antibody

Catalog No: #C43491

Package Size: #C43491-AF350 100ul #C43491-AF405 100ul #C43491-AF488 100ul

#C43491-AF555 100ul #C43491-AF594 100ul #C43491-AF647 100ul

#C43491-AF680 100ul #C43491-AF750 100ul #C43491-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	WFDC3 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification
Species Reactivity	Hu
Immunogen Description	Synthetic peptide of human WFDC3
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Target Name	WFDC3
Other Names	WAP14; dJ447F3.3
Accession No.	Swiss-Prot#: Q8IUB2 NCBI Gene ID: NP_542181
Uniprot	Q8IUB2
GeneID	140686;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at -20°C/1 year

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

This gene encodes a member of the WAP-type four-disulfide core (WFDC) domain family. The WFDC domain, or WAP signature motif, contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor. The encoded protein contains four WFDC domains. Most WFDC genes are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the telomeric cluster. Alternatively spliced transcript variants have been observed but their full-length nature has not been determined.

---

Note: This product is for in vitro research use only