Pyk2(Phospho-Tyr402) Antibody

Catalog No: #11216

Package Size: #11216-1 50ul #11216-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description				
Product Name	Pyk2(Phospho-Tyr402) Antibody			
Host Species	Rabbit			
Clonality	Polyclonal			
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.			
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho			
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.			
Applications	WB IHC IF			
Species Reactivity	Hu Ms Rt			
Specificity	The antibody detects endogenous level of Pyk2 only when phosphorylated at tyrosine 402.			
Immunogen Type	Peptide-KLH			
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 402 (D-I-Y(p)-A-E) derived from Human Pyk2.			
Target Name	Pyk2			
Modification	Phospho			
Other Names	FADK 2; FAK2; Focal adhesion kinase 2; PTK2B; Proline-rich tyrosine kinase 2 RAFTK			
Accession No.	Swiss-Prot: Q14289NCBI Protein: NP_004094.3			
Uniprot	Q14289			
GenelD	2185;			
Concentration	1.0mg/ml			
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%			
	sodium azide and 50% glycerol.			
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.			

Application Details			
Predicted MW: 116kd			
Western blotting: 1:500~1:1000			
Immunohistochemistry: 1:50~1:	00		
Immunofluorescence: 1:100~1:2	00		

Images



Western blot analysis of extracts from 293 cells untreated or treated with Serum using Pyk2(Phospho-Tyr402) Antibody #11216.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Pyk2(Phospho-Tyr402) Antibody #11216(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using Pyk2(Phospho-Tyr402) Antibody #11216.

Background

Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation. Gluck SL, et al. (2004) J Clin Invest; 114(12): 1696-1699 Benzing T, et al. (2001) Proc Natl Acad Sci U S A; 98(17): 9784-9789 Tian D, et al. (2002) Mol Cell Biol; 22(8): 2650-2662 Lu Z, et al. (2001) Mol Cell Biol; 21(12): 4016-4031

Krishnan HH, et al. (2006) J Virol; 80(3): 1167-1180

Note: This product is for in vitro research use only