

PARC Antibody

Catalog No: #24204




Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	PARC Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB ICC
Species Reactivity	Hu Ms
Immunogen Type	Peptide
Immunogen Description	Raised against a 17 amino acid peptide from near the carboxy terminus of human PARC.
Target Name	PARC
Other Names	p53-associated parkin-like cytoplasmic protein, Ubch7-associated protein 1
Accession No.	Swiss-Prot:Q8IWT3Gene ID:23113
Uniprot	Q8IWT3
GeneID	23113;
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of PARC in Daudi lysate with PARC antibody at (A) 1 and (B) 2 ug/mL.



Immunocytochemistry of Parc in Daudi cells with Parc antibody at 1 ug/mL.

Background

The continued localization of p53 to the nucleus is essential for its function as a tumor suppressor. PARC, a large, Parkin-like ubiquitin ligase has recently been identified as a cytoplasmic anchor protein in p53-associated protein complexes. In the absence of stress, PARC inactivation results in nuclear localization of p53 and activation of p53-dependent apoptosis, while overexpression of this protein promoted cytoplasmic sequestration of p53. Surprisingly, PARC knockout mice were viable and exhibited no obvious phenotype, suggesting that other proteins, such as the highly related cullin family of E3 ubiquitin ligases, may perform similar functions in the absence of PARC. Additionally, it has been suggested that p53 binding to PARC may serve to control PARC function.

Note: This product is for in vitro research use only