

## Rad17 (phospho-Ser635) rabbit pAb

Catalog No: #13589



Package Size: #13589-1 50ul #13589-2 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Rad17 (phospho-Ser635) rabbit pAb
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Applications	WB
Species Reactivity	Human
Specificity	This antibody detects endogenous levels of Human Rad17 (phospho-Ser635)
Immunogen Description	Synthesized phospho peptide around human Rad17 (Ser635)
Other Names	Cell cycle checkpoint protein RAD17 (hRad17) (RF-C/activator 1 homolog)
Accession No.	Swiss Prot:O75943GeneID:5884
Uniprot	O75943
GeneID	5884
SDS-PAGE MW	77
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

## Application Details

WB 1:1000-2000

## Background

RAD17 checkpoint clamp loader component(RAD17) Homo sapiens The protein encoded by this gene is highly similar to the gene product of *Schizosaccharomyces pombe* rad17, a cell cycle checkpoint gene required for cell cycle arrest and DNA damage repair in response to DNA damage. This protein shares strong similarity with DNA replication factor C (RFC), and can form a complex with RFCs. This protein binds to chromatin prior to DNA damage and is phosphorylated by the checkpoint kinase ATR following damage. This protein recruits the RAD1-RAD9-HUS1 checkpoint protein complex onto chromatin after DNA damage, which may be required for its phosphorylation. The phosphorylation of this protein is required for the DNA-damage-induced cell cycle G2 arrest, and is thought to be a critical early event during checkpoint signaling in DNA-damaged cells. Multiple alternatively spliced transcript variants of this gene, which encode four distinct protein isoforms, h

Note: This product is for in vitro research use only