## MARK1/2/3/4 (phospho Thr215) Polyclonal Antibody

Catalog No: #13731

SAB Signalway Antibody

Package Size: #13731-1 50ul #13731-2 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	MARK1/2/3/4 (phospho Thr215) Polyclonal Antibody
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific
	immunogen.
Applications	IHC-p,IF(paraffin section),ELISA
Species Reactivity	Human,Mouse,Rat
Specificity	Phospho-MARK1/2/3/4 (T215) Polyclonal Antibody detects endogenous levels of MARK1/2/3/4 protein only
	when phosphorylated at T215.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human MARK1/2/3/4 around the
	phosphorylation site of Thr215. AA range:181-230
Other Names	MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating
	kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL
	motif kinase 1; EMK-1; MAP/microtubule affin
Accession No.	Swiss Prot:Q9P0L2/Q7KZI7/P27448/Q96L34GeneID:4139/2011/4140/57787
Uniprot	Q9P0L2/Q7KZI7/P27448/Q96L34
GeneID	4139/2011/4140/57787
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

## **Application Details**

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

## Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Thr-215 by STK11 in complex with STE20-related adapter-alpha (STRAD alpha) pseudo kinase and CAB39.,function:May play a role in cytoskeletal stability.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. MARK subfamily.,similarity:Contains 1 KA1 (kinase-associated) domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UBA domain.,subcellular location:Appears to localize to an intracellular network.,tissue specificity:Highly expressed in heart, skeletal muscle, brain, fetal brain and fetal kidney.,

Note: This product is for in vitro research use only