

c-Cbl (phospho-Tyr731) rabbit pAb

Catalog No: #13996

Package Size: #13996-1 50ul #13996-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	c-Cbl (phospho-Tyr731) rabbit pAb
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Applications	WB
Species Reactivity	Human
Specificity	This antibody detects endogenous levels of Human c-Cbl (phospho-Tyr731)
Immunogen Description	Synthesized phosho peptide around human c-Cbl (Tyr731)
Other Names	E3 ubiquitin-protein ligase CBL (EC 6.3.2.-) (Casitas B-lineage lymphoma proto-oncogene) (Proto-oncogene c-Cbl) (RING finger protein 55) (Signal transduction protein CBL)
Accession No.	Swiss Prot:P22681GenelD:867
Uniprot	P22681
GenelD	867
SDS-PAGE MW	100
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

Application Details

WB 1:1000-2000

Background

Cbl proto-oncogene(CBL) Homo sapiens This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5' UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause of Noonan syndrome-like disorder. [provided by RefSeq, Jul 2016].

Note: This product is for in vitro research use only