Recombinant human IL15

Catalog No: #AG0011

Description



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	Recombinant human IL15
Host Species	HEK293
Purification	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC
Immunogen Description	Asn49-Ser162
Target Name	IL15
Other Names	Human IL-15, h-IL-15, rh-IL-15, recombinant IL-15, interleukin-15
Accession No.	Uniprot:P40933Gene ID:3600
Uniprot	P40933
GeneID	3600
Target Species	human
Calculated MW	12.8 KDa
Tag Info	addtional amino acid free
Formulation	0.22 μm filtered solution of PBS, pH 7.4.
Storage	Aliquot and store at -80°C. Avoid repeated freeze/thaw cycles.

Background

Interleukin 15 (IL-15) is a widely expressed 14?kDa cytokine that is structurally and functionally related to IL-2 and plays an important role in many immunological diseases (1, 2). Mature human IL-15 protein shares 70% amino acid sequence identity with mouse and rat IL-15. Alternative splicing generates isoforms of Interleukin 15 with either a long or short signal peptide (LSP or SSP), and the SSP isoform is retained intracellularly (3). The IL-15 protein binds with high affinity to IL-15 R alpha (4). It binds with lower affinity to a complex of IL-2 R beta and the common gamma chain?(gamma c) which are also subunits of the IL-2 receptor complex (5). IL-15 associates with IL-15 R alpha in the endoplasmic reticulum, and this complex is expressed on the cell surface (6). The dominant mechanism of IL-15 action is known as transpresentation in which IL-15 and IL-15 R alpha are coordinately expressed on the surface of one cell and interact with complexes of IL-2 R beta / gamma c on adjacent cells (7). This enables cells to respond to Interleukin 15 even if they do not express IL-15 R alpha (6). In human and mouse, soluble IL-15 R alpha functions as an inhibitor that limits IL-15 action (4,?9). Ligation of membrane-associated IL-15/IL-15 R alpha complexes also induces reverse signaling that promotes activation of the IL-15/IL-15 R alpha expressing cells (11). IL-15 induces or enhances the differentiation, maintenance, or activation of multiple T?cell subsets including NK, NKT, Th17, Treg, and CD8+?memory cells (12 - 16). An important component of these functions is the ability of?IL-15 to induce dendritic cell differentiation and inflammatory activation (11, 14). IL-15 exhibits anti-tumor activity independent of its actions on NK cells or CD8+?T?cells (17). It also inhibits the deposition of lipid in adipocytes, and its circulating levels are decreased in obesity (18).

Note: This product is for in vitro research use only