

Recombinant mouse IL17A

Catalog No: #AG0042

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Recombinant mouse IL17A
Host Species	HEK293
Purification	> 95% by Tris-Bis PAGE; > 95% by SEC-HPLC
Immunogen Description	Ala26-Ala158
Target Name	IL17A
Other Names	CTLA8; CTLA-8; CTLA8cytotoxic T-lymphocyte-associated serine esterase 8; Cytotoxic T-lymphocyte-associated antigen 8; IL17; IL-17; IL17A; IL-17A; IL-17Acytotoxic T-lymphocyte-associated protein 8; IL-17CTLA-8; IL17interleukin-17A; interleukin 17 (cytotoxic T-lymphocyte-associated serine esterase 8); interleukin 17A
Accession No.	Uniprot:Q62386Gene ID:16171
Uniprot	Q62386
GeneID	16171
Target Species	mouse
Calculated MW	15.0 KDa
Tag Info	additional amino acid free
Formulation	0.22 µm filtered solution of PBS, pH 7.4.
Storage	Aliquot and store at -80°C. Avoid repeated freeze/thaw cycles.

Background

Interleukin-17A (IL-17A), also known as CTLA-8, is a 15-20 kDa glycosylated cytokine that plays an important role in anti-microbial and chronic inflammation. The six IL-17 cytokines (IL-17A-F) are encoded by separate genes but adopt a conserved cystine knot fold (1, 2). Mature mouse IL-17A shares 61% and 89% amino acid sequence identity with human and rat IL-17A, respectively (3, 4). IL-17A is secreted by Th17 cells, gamma /P' T cells, iNKT cells, NK cells, LT α i cells, neutrophils, and intestinal Paneth cells (2). It forms disulfide-linked homodimers as well as disulfide-linked heterodimers with IL-17F (5, 6). IL-17A exerts its effects through the transmembrane IL-17RA in complex with IL-17RC or IL-17RD (7, 8). Both IL-17RA and IL-17RC are required for responsiveness to heterodimeric IL-17A/F (7). IL-17A promotes protective mucosal and epidermal inflammation in response to microbial infection (9-12). It induces chemokine production, neutrophil influx, and the production of antibacterial peptides (9-11). IL-17A/F likewise induces neutrophil migration, but IL-17F does not (11). IL-17A additionally enhances the production of inflammatory mediators by rheumatoid synovial fibroblasts and contributes to TNF-alpha induced shock (Fossiez, 14). In contrast, it can protect against the progression of colitis by limiting chronic inflammation (12). IL-17A encourages the formation of autoreactive germinal centers and exacerbates the onset and progression of experimental models of autoimmunity (15, 16). IL-17A has been shown to exert either tumorigenic or anti-tumor effects (17, 18).

Note: This product is for in vitro research use only