

Recombinant mouse FLT3L

Catalog No: #AG0046

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Recombinant mouse FLT3L
Host Species	HEK293
Purification	> 95% by Tris-Bis PAGE; > 95% by SEC-HPLC
Immunogen Description	Gly27-Arg188
Target Name	FLT3L
Other Names	FL; FLG3L; Flt3 ligand; Flt-3 Ligand; Flt3L; FLT3LG; fms-related tyrosine kinase 3 ligand; SL cytokine
Accession No.	Uniprot:P49772Gene ID:14256
Uniprot	P49772
GeneID	14256
Target Species	mouse
Calculated MW	18.4 KDa
Tag Info	additional amino acid free
Formulation	0.22 µm filtered solution of PBS, pH 7.4.
Storage	Aliquot and store at -80°C. Avoid repeated freeze/thaw cycles.

Background

Flt-3 Ligand, also known as FLT3L, is an alpha-helical cytokine that promotes the differentiation of multiple hematopoietic cell lineages (1-3). Mature mouse Flt-3 Ligand consists of a 161 amino acid (aa) extracellular domain (ECD) with a cytokine-like domain and a juxtamembrane tether region, a 21 aa transmembrane segment, and a 22 aa cytoplasmic tail (4-6). Within the ECD, mouse Flt-3 Ligand shares 71% and 81% aa sequence identity with human and rat Flt-3 Ligand, respectively. The mouse and human Flt-3 Ligand proteins show cross-species activity (4, 5, 7). Flt-3 Ligand is also structurally related to M-CSF and SCF. Flt-3 Ligand is widely expressed in various mouse and human tissues. Flt-3 Ligand is expressed as a noncovalently-linked dimer by T cells and bone marrow and thymic fibroblasts (1, 8). Each 36 kDa chain of the Flt-3 Ligand dimer carries approximately 12 kDa of N- and O-linked carbohydrates (8). Alternate splicing and proteolytic cleavage of the transmembrane form of the Flt-3 Ligand protein can generate a soluble 30 kDa fragment that includes the cytokine-like domain (4, 8). Alternate splicing of mouse Flt-3 Ligand also generates a membrane-associated isoform with a 57 aa substitution following the cytokine-like domain in the ECD of the Flt-3 Ligand protein (4, 5, 8, 9). Both transmembrane and soluble Flt-3 Ligand signal through the tyrosine kinase receptor Flt-3/Flk-2 (3 - 6). Flt-3 Ligand induces the expansion of monocytes and immature dendritic cells as well as early B cell lineage differentiation (2, 10). Additionally, Flt-3 Ligand synergizes with IL-3, GM-CSF, and SCF to promote the mobilization and myeloid differentiation of hematopoietic stem cells (4, 5, 7). Flt-3 Ligand also cooperates with IL-2, IL-6, IL-7, and IL-15 to induce NK cell development and with IL-3, IL-7, and IL-11 to induce terminal B cell maturation (1, 11). Animal studies also show that Flt-3 Ligand reduces the severity of experimentally induced allergic inflammation (12).

Note: This product is for in vitro research use only