OCIAD1 Antibody

Catalog No: #24995

Description



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

OCIAD1 Antibody
Rabbit
Polyclonal
Affinity chromatography purified via peptide column
ELISA WB
Hu
Peptide
Raised against a 16 amino acid peptide from near the carboxy terminus of human OCIAD1.
OCIAD1
OCIAD1, Ovarian cancer immuno-reactive antigen domain-containing protein 1, OCIA domain-containing
protein 1
Swiss-Prot:Q9NX40Gene ID:54940
Q9NX40
54940;
1mg/ml
Supplied in PBS containing 0.02% sodium azide.
Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated
freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of OCIAD1 in 293 cell lysate with OCIAD1 antibody at (A) 1 and (B) 2 ug/mL.

Background

OCIAD1 was identified via immunoscreening of an ovarian carcinoma cDNA library from ovarian cancer patients and is expressed in multiple tissues including ovary, placenta, brain, testis, prostate, and mammary gland. Two isoforms of OCIAD1 are known to exist; the shorter isoform is restricted to the central nervous system. OCIAD1 is a transmembrane protein whose overexpression in HEY ovarian cancer cells increased lysophosphatidic acid-(LPA-)induced, but not basal level cell adhesion to extracellular matrix proteins collagen I and laminin10/11. This adhesion is not blocked by LY294002 and GF109203X, suggesting that OCIAD1 does not use protein kinase C and PI3 kinase signaling pathways to exert its effect on adhesion.

Note: This product is for in vitro research use only