JNK1/JNK2/JNK3(phospho-Thr183/Tyr185) Antibody

Catalog No: #11504

Package Size: #11504-1 50ul #11504-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description					
Product Name	JNK1/JNK2/JNK3(phospho-Thr183/Tyr185) Antibody				
Host Species	Rabbit				
Clonality	Polyclonal				
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.				
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho				
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.				
Applications	WB IF				
Species Reactivity	Hu Ms Rt				
Specificity	The antibody detects endogenous level of JNK1/JNK2/JNK3 only when phosphorylated at Thr183/Tyr185.				
Immunogen Type	Peptide-KLH				
Immunogen Description	Peptide sequence around phosphorylation site of Thr183/Tyr185 (M-M-T(p)-P-Y(p)- V - V) derived from				
	Human JNK1/JNK2/JNK3.				
Target Name	JNK1/JNK2/JNK3				
Modification	Phospho				
Other Names	Stress-activated protein kinase JNK1; c-Jun N-terminal kinase 1; JNK-46				
Accession No.	Swiss-Prot: P45983 P45984 P53779NCBI Protein: NP_002741.1 NP_001128516.1 NP_002744.1				
Uniprot	P45983				
GenelD	5599;				
Concentration	1.0mg/ml				
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%				
	sodium azide and 50% glycerol.				
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.				

Predicted MW: 46 54 kd Western blotting: 1:500~1:1000	Application Details			
	Predicted MW: 46 54 kd			
Immunofluorescence: 1:100~1:200	Western blotting: 1:500~1:1000	D		
	Immunofluorescence: 1:100~1	:200		


Images



Western blot analysis of extracts from C6 cells untreated or treated with anisomycin using JNK1/JNK2/JNK3(phospho-Thr183/Tyr185) Antibody #11504.



Immunofluorescence staining of methanol-fixed Hela cells using JNK1/JNK2/JNK3(phospho-Thr183/Tyr185) Antibody #11504.




PC12

Cd NAC

Neuron

Cd+59600125

Western blot analysis of extracts from 293 cells, treated with Anisomycin or calf intestinal phosphatase (CIP), using JNK1/JNK2/JNK3 (phospho-Thr183/Tyr185) Antibody #11504.



Background

kDa 55

40

Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells By similarity. Phosphorylates heat shock factor protein 4 (HSF4). /Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required differentiation of T-helper cells into Th1 cells. JNK2 isoforms display different binding patterns: a-1 and a-2 preferentially bind to c-Jun, whereas beta-1 and beta-2 bind to ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms. JUNB is not a

JNK (p-Thr183/Tyr185)

substrate for JNK2 a-2, and JUND binds only weakly to it. /Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. Required for stress-induced neuronal apoptosis and the pathogenesis of glutamate excitotoxicity Davis, R.J. (1999) Biochem Soc Symp 64, 1-12.

Ichijo, H. (1999) Oncogene 18, 6087-93.

Kyriakis, J.M. and Avruch, J. (2001) Physiol Rev 81, 807-69.

Note: This product is for in vitro research use only