## Rad50 Rabbit mAb

Catalog No: #52037

Package Size: #52037-1 50ul #52037-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_					
	es	$\circ$ r	nt	-	n
		C I I	IUI	IU	

Product Name	Rad50 Rabbit mAb	
Host Species	Recombinant Rabbit	
Clonality	Monoclonal antibody	
Clone No.	S05-5H6	
Isotype	Rabbit IgG	
Purification	Affinity Purified	
Applications	WB	
Species Reactivity	Human, Mouse	
Immunogen Description	A synthetic peptide of human Rad50	
Conjugates	Unconjugated	
Modification	Unmodification	
Other Names	NBSLD; RAD502; hRad50	
Accession No.	Swiss-Prot:Q92878GeneID:10111	
Uniprot	Q92878	
GeneID	10111	
Calculated MW	Calculated MW: 154 kDa; Observed MW: 154 kDa	
Concentration	0.3 mg/ml	
Formulation	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.	

## Application Details

WB: 1/2000

## **Images**



Western blot detection of Rad50 in K562,Hela cell lysates using Rad50 Rabbit mAb(1:1000 diluted).Predicted band size:154kDa.Observed band size:154kDa.

## Background

Swiss-Prot Acc.Q92878.Component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis. The complex possesses single-strand endonuclease activity and double-strand-specific 3'-5' exonuclease activity, which are provided by MRE11. RAD50 may be required to bind DNA ends and hold them in close proximity. This could facilitate searches for short or long regions of sequence homology in the recombining DNA templates, and may also stimulate the activity of DNA ligases and/or restrict the nuclease activity of MRE11 to prevent nucleolytic degradation past a given point (PubMed:11741547, PubMed:9590181, PubMed:9705271, PubMed:9651580).

The complex may also be required for DNA damage signaling via activation of the ATM kinase (PubMed:15064416).

In telomeres the MRN complex may modulate t-loop formation (PubMed:10888888).

Note: This product is for in vitro research use only