CDKN2A Rabbit mAb

Catalog No: #52214

Package Size: #52214-1 50ul #52214-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | |
|-------------------|-------|-----|-----|
| | escri | nti | n |
| $\boldsymbol{ u}$ | COUL | μu | ULI |

| Product Name | CDKN2A Rabbit mAb | |
|-----------------------|--|--|
| Host Species | Recombinant Rabbit | |
| Clonality | Monoclonal antibody | |
| Clone No. | S04-8C9 | |
| Isotype | Rabbit IgG | |
| Purification | Affinity Purified | |
| Applications | WB IF | |
| Species Reactivity | Human, Mouse | |
| Immunogen Description | Recombinant protein of human CDKN2A/p16INK4a | |
| Conjugates | Unconjugated | |
| Modification | Unmodification | |
| Other Names | ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; | |
| | P16INK4; P16INK4A; P16-INK4A | |
| Accession No. | Swiss-Prot:P42771GeneID:1029 | |
| Uniprot | P42771 | |
| GeneID | 1029 | |
| Calculated MW | Calculated MW: 17 kDa; Observed MW: 17 kDa | |
| Concentration | 0.3 mg/ml | |
| Formulation | 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA | |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. | |


Application Details

WB: 1/1000; ICC/IF: 1/20-1/100;


Images



Western blot detection of CDKN2A/p16INK4a in 293 cell lysates using CDKN2A/p16INK4a Rabbit mAb(1:1000 diluted). Predicted band size:17KDa. Observed band size:17KDa.



Western blot detection of CDKN2A-p16INK4a in 3T3,Hela cell lysates using CDKN2A/p16INK4a Rabbit mAb(1:1000 diluted).Predicted band size:17KDa.Observed band size:17KDa.



Immunofluorescence of CDKN2A/p16INK4a(green) in Hela cells using CDKN2A/p16INK4a Rabbit mAb at dilution 1/5, and DAPI(blue)

Background

Swiss-Prot Acc.P42771.Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

Note: This product is for in vitro research use only