

IDH1 Antibody


Catalog No: #25141

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	IDH1 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide
Immunogen Description	Raised against a 13 amino acid peptide near the carboxy terminus of human IDH1.
Target Name	IDH1
Other Names	Oxalosuccinate decarboxylase, Cytosolic NADP-isocitrate dehydrogenase, NADP(+)-specific ICDH, IDCD, IDH, IDP, PICD
Accession No.	Swiss-Prot:O75874Gene ID:3417
Uniprot	O75874
GeneID	3417;
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of IDH1 in HepG2 cell lysate with IDH1 antibody at (A) 1 and (B) 2 ug/mL.

Background

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Two NADP(+)-dependent isocitrate dehydrogenases have been found as homodimer: IDH1 is predominantly cytosolic and peroxisomal and IDH2 is mitochondrial. The presence of IDH1 in peroxisomes suggests it may play a role in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic IDH1 serves a significant role in cytoplasmic NADPH production. Defects in IDH1 are involved in the development of glioma.

Note: This product is for in vitro research use only