CDKN2A/p14ARF Rabbit mAb

Catalog No: #56368

Package Size: #56368-1 50ul #56368-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | | |
|---|----|-----|-----|-----|
| П | 00 | ori | nti | ion |
| U | ヒ٥ | UH | บแ | UI |

| Product Name | CDVN2A/p14ARE Robbit mAb |
|-----------------------|--|
| FIGURE INAME | CDKN2A/p14ARF Rabbit mAb |
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB ICC/IF IP FC |
| Species Reactivity | Human |
| Specificity | CDKN2A/p14ARF Antibody detects endogenous levels of total CDKN2A/p14ARF |
| Immunogen Description | A synthesized peptide derived from human CDKN2A/p14ARF |
| Other Names | ARF; CDK4I; CDKN2; CDKN2A; CMM2; INK4; INK4a; MLM; MTS1; p14; p16; p16INK4a; P19ARF; TP16; |
| Accession No. | Uniprot:Q8N726 |
| Uniprot | Q8N726 |
| Calculated MW | 14kDa |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at +4Λ C short term. Store at -20Λ C long term. Avoid freeze / thaw cycle. |

Application Details

WB:1:500~1:1000ICC/IF:1:50~1:200IP:1:50FC:1:80

Images



Western blot analysis of CDKN2A/p14ARF expression in PC3 cell lysate.

Product Description

The gene for CDK2NA generates several transcripts/proteins which differ from each other in their first exons. Three of these transcripts are generated by alternative splicing (isoform 1 a.k.a p16INK4A, isoform 2 and isoform 3 a.k.a p12), two of which are known to function as inhibitors of CDK4 kinase.

Note: This product is for in vitro research use only