## HIF Prolyl Hydroxylases Rabbit mAb

Catalog No: #57046

Package Size: #57046-1 50ul #57046-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$		4.0	
	escri	ntı	nη
$\boldsymbol{ u}$	COUL	Μu	UH

Product Name	HIF Prolyl Hydroxylases Rabbit mAb	
Host Species	Rabbit	
Clonality	Monoclonal	
Isotype	Rabbit IgG	
Purification	Affinity-chromatography	
Applications	WB IHC IP FC	
Species Reactivity	Human	
Specificity	HIF Prolyl Hydroxylases Antibody detects endogenous levels of total HIF Prolyl Hydroxylases	
Immunogen Description	A synthesized peptide derived from human HIF Prolyl Hydroxylases	
Other Names	EGLN4; HIFPH4; Hypoxia inducible factor prolyl 4 hydroxylase; P4H with transmembrane domain; P4htm;	
	PH4; PHD4; Proline 4 hydroxylase; Prolyl hydroxlase domain containing 4;	
Accession No.	Uniprot:Q9NXG6	
Uniprot	Q9NXG6	
Calculated MW	47kDa	
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	
Storage	Store at +4Λ C short term. Store at -20Λ C long term. Avoid freeze / thaw cycle.	

## **Application Details**

WB:1:500~1:2000IHC:1:50~1:200IP:1:50FC:1:50

## **Images**



Western blot analysis of HIF Prolyl Hydroxylases expression in HeLa cell lysate.

## **Product Description**

Catalyzes the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates HIF1A at 'Pro-402' and 'Pro-564'. May function as a cellular oxygen sensor and, under normoxic conditions, may target HIF through the hydroxylation for proteasomal degradation via the von Hippel-Lindau ubiquitination complex.

Note: This product is for in vitro research use only