

TrkA Antibody FITC Conjugated

Catalog No: #C00651F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	TrkA Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human TrkA
Conjugates	FITC
Target Name	TrkA
Other Names	CIPA; CIPA; DKFZp781I14186; gp140trk; High affinity nerve growth factor receptor; High affinity nerve growth factor receptor precursor; High affinity nerve growth factor receptor precursor; MTC; MTC; neurotrophic tyrosine kinase receptor type 1; neurotrophic tyrosine kinase receptor type 1; Ntrk 1;
Accession No.	NCBI Gene ID4914
Uniprot	P04629
GeneID	4914;
Excitation Emission	494nm 518nm
Cell Localization	Extracellular
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

ICC=1:50-200 IF=1:50-200

Background

The Trk family of nerve growth factor receptors includes Trk A(also referred to as Trk A gp140), Trk B and Trk C. The prototype member of this gene family, Trk A, encodes a 140 kDa cell surface receptor , gp140, the expression of which is restricted in vivo to neurons of the sensory spinal and cranial ganglia of neurocrest origin. Nerve growth factor (NGF) stimulates tyrosine phosphorylation of Trk gp 140 in neural cell lines and in embryonic dorsal root ganglia. By comparison, BDNF and to a lesser extent, NT-3, but not NGF, can induce tyrosine phosphorylation of Trk B gp 145. The third member of the Trk receptor family, Trk C encodes a 140 kDa protein, Trk C gp140, that is preferentially expressed in brain tissue and primarily functions as a receptor for NT-3. An additional component of the Trk receptor complex, NGFR p175, binds to neurotrophic factors with low affinity but is required for efficient signaling. NGFR p175 accelerates Trk activation and may recruit downstream effector molecules to the ligand-bound receptor complex.

Note: This product is for in vitro research use only