## MHC class I Antibody FITC Conjugated

Catalog No: #C00670F

Description



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	MHC class I Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 60-105 181 derived from human MHC class I
Conjugates	FITC
Target Name	MHC class I
Other Names	A 28; A 9; Aw 24; Aw 68; CLASS I HISTOCOMPATIBILITY ANTIGEN; H2 K1; H2K; HLA A; HLA-A; HLA C;
	HLA-C; HLA class I histocompatibility antigen A 1 alpha chain; HLA class I histocompatibility antigen A 11
	alpha chain; HLA class I histocompatibility antigen A 2 alpha chain; HLA class I histocompatibility
Accession No.	Swiss-Prot#G0ZMJ8
Uniprot	G0ZMJ8
Excitation Emission	494nm 518nm
Cell Localization	Extracellular
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## **Application Details**

ICC=1:50-200 IF=1:50-200

## Background

HLA-A belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon 1 encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domains, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Hundreds of HLA-A alleles have been described. [provided by RefSeq, Jul 2008]

Note: This product is for in vitro research use only