USP11 Rabbit mAb

Catalog No: #59585

Package Size: #59585-1 50ul #59585-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | | | |
|------------------|----|-----|----|----|-----|
| רו | es | Cri | nt | ī | n |
| \boldsymbol{L} | しつ | UH | νι | IU | 4 8 |

| Product Name | USP11 Rabbit mAb | |
|-----------------------|--|--|
| Host Species | Rabbit | |
| Clonality | Monoclonal | |
| Isotype | Rabbit IgG | |
| Purification | Affinity-chromatography | |
| Applications | WB ICC/IF IP FC | |
| Species Reactivity | Human Mouse Rat | |
| Specificity | USP11 Antibody detects endogenous levels of total USP11 | |
| Immunogen Description | A synthesized peptide derived from human USP11 | |
| Other Names | UHX1; USP11; | |
| Accession No. | Uniprot:P51784 | |
| Uniprot | P51784 | |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. | |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. | |

Application Details

WB 1:500~1:2000 ICC/IF 1:50~1:200 IP 1:100 FC 1:200

Images



Western blot analysis of USP11 expression in (1) HEK293 cell lysate; (2) Mouse testis lysate.

Product Description

Protease that can remove conjugated ubiquitin from target proteins and polyubiquitin chains. Inhibits the degradation of target proteins by the proteasome. Plays a role in the regulation of pathways leading to NF-kappa-B activation. Plays a role in the regulation of DNA repair after double-stranded DNA breaks.

Background

Protease that can remove conjugated ubiquitin from target proteins and polyubiquitin chains. Inhibits the degradation of target proteins by the proteasome. Plays a role in the regulation of pathways leading to NF-kappa-B activation. Plays a role in the regulation of DNA repair after double-stranded DNA breaks.

Note: This product is for in vitro research use only