CD45 Antibody FITC Conjugated

Catalog No: #C00815F



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Support: tech@signalwayantibody.con
CD45 Antibody FITC Conjugated
Rabbit
Polyclonal
IgG
Purified by Protein A.
Flow-Cyt,ICC,IF
Hu Ms Rt
KLH conjugated synthetic peptide aa 500-530 1291 derived from mouse CD45
FITC
CD45
Receptor-type tyrosine-protein phosphatase C, Leukocyte common antigen, Lymphocyte antigen 5, L-CA,
Ly-5, T200
Swiss-Prot#P06800
P06800
19264;
494nm 518nm
Extracellular
1mg ml
0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=1ug/Test ICC=1:50-200 IF=1:50-200

Background

Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN (By similarity). Dephosphorylates LYN, and thereby modulates LYN activity.

Note: This product is for in vitro research use only