

TLX2 Antibody FITC Conjugated

Catalog No: #C01703F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	TLX2 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	ICC IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human TLX2
Conjugates	FITC
Target Name	TLX2
Other Names	Enx; homeo box 11 like 1; Homeobox protein Hox 11L1; Homeobox TLX 2; HOX11L1; NCX; Neural crest homeobox protein; PMUR10F; T cell leukemia homeobox protein 2; T cell leukemia, homeobox 2.
Excitation Emission	494nm 518nm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

ICC=1:50-200 IF=1:50-200

Background

T-cell leukemia homeobox protein 2 (TLX2), also known as homeobox protein Hox-11L1 (HOX11L1), neural crest homeobox protein (NCX) or ENX, is a 284 amino acid member of the TLX homeobox family. The mouse homolog, Tlx2, has been detected in dorsal-root ganglia, cranial and enteric-nerve ganglia, parasympathetic ganglia and adrenal glands in mouse embryos and in the adrenal glands, intestine and heart of adult mice. The expression pattern of Tlx2, which is restricted to tissues derived from neural crest cells, suggests that it may play a role in the proliferation or differentiation of the enteric peripheral nervous system. TLX2, which is localized to the nucleus, is highly homologous to mouse Tlx2, and shares several critical domains, including an enhancer element in the promoter that is crucial for tissue-specific expression. Mutations in the gene encoding mouse Tlx2 lead to congenital anomalies closely resembling neuronal intestinal dysplasia in humans. Thus, TLX2 is thought to play a role in this disease, which is a rare condition characterized by hyperplasia of submucosal plexus with giant submucosal ganglia and increased acetylcholinesterase activity in nerve fiber around submucosal blood vessels.

Note: This product is for in vitro research use only