

BOB1 Antibody HRP Conjugated

Catalog No: #C02159H

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	BOB1 Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	WB IHC-P IHC-F ICC
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide derived from human BOB1
Conjugates	HRP
Target Name	BOB1
Other Names	B cell Oct binding protein 1; B cell specic coactivator OBF 1; B cell specic coactivator OBF1; B-cell-specic coactivator OBF-1; BOB 1; BOB-1; OBF 1; OBF1; OBF1_HUMAN; OCA B; OCA-B; OCAB; OCT binding factor 1; OCT-binding factor 1; POU domain class 2 associating factor 1; POU domain class 2-associati
Accession No.	NCBI Gene ID5450
Uniprot	Q16633
GeneID	5450;
Excitation Emission	N A
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

WB=1:500-2000 IHC-P=1:50-200 IHC-F=1:50-200 ICC=1:50-200

Background

POU domain proteins contain a bipartite DNA-binding domain divided by a flexible linker that enables them to adopt various monomer configurations on DNA. The versatility of POU protein operation is additionally conferred at the dimerization level. The POU dimer from the OCT1 gene formed on the palindromic OCT factor recognition element, or PORE (ATTTGAAATGCAAAT), could recruit the transcriptional coactivator OBF1. Studies of tissue-specific expression of immunoglobulin promoters demonstrate the importance of an octamer, ATTTGCAT, and the proteins that bind to it. This is a regulatory element important for tissue- and cell-specific transcription as well as for transcription of a number of housekeeping genes. Oct-1 encodes one protein, NF-A1, which is found in nuclear extracts from all cell types and thus is not specific to lymphoid cells as is the protein NF-A2, which is encoded by Oct-2. A novel protein designated Bob 1 (B cell Oct binding protein 1), alternatively called OBF-1, specifically interacts with Oct-1 and Oct-2, enhancing their transcriptional efficacy. Bob 1 is expressed at highest levels in spleen and peripheral blood leukocytes and represents an Oct co-factor capable of conferring cell-specific activation of Oct-1 and Oct-2. Although having no intrinsic capacity for DNA binding, Bob 1 associates tightly with the octamer motif in the presence of Oct-1 and or Oct-2. The gene which encodes Bob 1 maps to human chromosome 11q23.1.

Note: This product is for in vitro research use only