Akt(Ab-473) Antibody

Catalog No: #21054

Package Size: #21054-1 50ul #21054-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


| the state of the s | |
|--|---|
| Product Name | Akt(Ab-473) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were |
| | purified by affinity-chromatography using epitope-specific peptide. |
| Applications | WB IHC IF |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous level of total Akt protein. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around aa. 471~475 (Q-F-S-Y-S) derived from Human Akt. |
| Target Name | Akt |
| Other Names | AKT; C-AKT; PKB; PKB-alpha; RAC |
| Accession No. | Swiss-Prot: P31749NCBI Protein: NP_001014431.1 |
| Uniprot | P31749 |
| GeneID | 207; |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |
| | |

Application Details


Predicted MW: 60kd

Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100


Images



Western blot analysis of extracts from 293 and 3T3 cells using Akt(Ab-473) Antibody #21054.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Akt(Ab-473) Antibody #21054(left) or the same antibody preincubated with blocking peptide(right).



Western blot analysis of extracts of various cell lines, using Akt(Ab-473) #21054

Background

General protein kinase capable of phosphorylating several known proteins. Phosphorylates TBC1D4. Signals downstream of phosphatidylinositol 3-kinase (PI3K) to mediate the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor I (IGF-I). Plays a role in glucose transport by mediating insulin-induced translocation of the GLUT4 glucose transporter to the cell surface. Mediates the antiapoptotic effects of IGF-I. Mediates insulin-stimulated protein synthesis by phosphorylating TSC2 at 'Ser-939' and 'Thr-1462', thereby activating mTORC1 signaling and leading to both phosphorylation of 4E-BP1 and in activation of RPS6KB1. Promotes glycogen synthesis by mediating the insulin-induced activation of glycogen synthase.

Baudhuin LM, et al. (2004) FASEB J Feb; 18(2): 341-3.

Min YH, et al. (2004) Cancer Res; 64(15): 5225-31.

Feng J, et al. (2004) J Biol Chem; 279(34): 35510-7.

Ayala G, et al. (2004) Clin Cancer Res; 10(19): 6572-8.

Lungu AO, et al. (2004) J Biol Chem; 279(47): 48794-800.

Note: This product is for in vitro research use only