

# Tap1 Antibody FITC Conjugated

Catalog No: #C04321F

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Tap1 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt IF
Species Reactivity	HuB MsB RtB B B B
Immunogen Description	KLH conjugated synthetic peptide aa 525-575 808 derived from human Tap1 ABCB2
Conjugates	FITC
Target Name	Tap1
Other Names	APT1; PSF1; ABCB2; PSF-1; RING4; TAP1N; D6S114E; TAP1*12N; Antigen peptide transporter 1; ATP-binding cassette sub-family B member 2; Peptide supply factor 1; Peptide transporter PSF1; Peptide transporter TAP1; Peptide transporter involved in antigen processing 1; Really interesting new gene
Accession No.	Swiss-Prot#Q03518NCBI Gene ID6890
Uniprot	Q03518
GeneID	6890;
Excitation Emission	494nm 518nm
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

## Application Details

Flow-Cyt=1:50-200B IF=1:50-200B

## Background

Involved in the transport of antigens from the cytoplasm to the endoplasmic reticulum for association with MHC class I molecules. Also acts as a molecular scaffold for the final stage of MHC class I folding, namely the binding of peptide. Nascent MHC class I molecules associate with TAP via tapasin. Inhibited by the covalent attachment of herpes simplex virus ICP47 protein, which blocks the peptide-binding site of TAP. Inhibited by human cytomegalovirus US6 glycoprotein, which binds to the luminal side of the TAP complex and inhibits peptide translocation by specifically blocking ATP-binding to TAP1 and prevents the conformational rearrangement of TAP induced by peptide binding. Inhibited by human adenovirus E3-19K glycoprotein, which binds the TAP complex and acts as a tapasin inhibitor, preventing MHC class I TAP association. Expression of TAP1 is down-regulated by human Epstein-Barr virus vL-1 protein, thereby affecting the transport of peptides into the endoplasmic reticulum and subsequent peptide loading by MHC class I molecules.

Note: This product is for in vitro research use only