ICAM-1(Ab-512) Antibody

Catalog No: #21106

Package Size: #21106-1 50ul #21106-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_		4.5	
Des	cri	ntic	าท
レしる	UH	ρ u	ווע


Product Name	ICAM-1(Ab-512) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB	
Species Reactivity	Hu	
Specificity	The antibody detects endogenous level of total ICAM-1 protein.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around aa. 510~514 (K-K-Y-R-L) derived from Human ICAM-1.	
Target Name	ICAM-1	
Other Names	ICA1; ICAM1;	
Accession No.	Swiss-Prot: P05362NCBI Protein: NP_000192.2	
Uniprot	P05362	
GeneID	3383;	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

Application Details

Predicted MW: 89 92kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from MDA cells using ICAM-1(Ab-512) Antibody #21106.

Background

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin a-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through SGEF and RHOG activation. In case of rhinovirus infection acts as a cellular receptor for the virus.

Greenwood J, et al. (2003) J Immunol; 171(4):2099-2108. Zhou Z, et al. (2005) Eur J Pharmacol; 513(1-2):1-8. Chen YH, et al. (2001) J Cell Biochem; 82(3):512-521

Note: This product is for in vitro research use only