

CYLD (Phospho-Ser418) Antibody HRP Conjugated

Catalog No: #C04469H

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	CYLD (Phospho-Ser418) Antibody HRP Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	WBB B IHC-PB IHC-F
Species Reactivity	HuB MsB RtB B
Immunogen Description	KLH conjugated synthetic phosphopeptide aa 390-440 956 derived from human CYLD around the phosphorylation site of Ser418
Conjugates	HRP
Target Name	CYLD Ser418
Other Names	EAC; MFT; SBS; TEM; BRSS; CDMT; MFT1; CYLD1; CYLDI; USPL2; Ubiquitin carboxyl-terminal hydrolase CYLD; Deubiquitinating enzyme CYLD; Ubiquitin thioesterase CYLD; Ubiquitin-specific-processing protease CYLD; CYLD; KIAA0849; HSPC057
Accession No.	Swiss-Prot#Q9NQC7NCBI Gene ID1540
Uniprot	Q9NQC7
GeneID	1540;
Excitation Emission	N A
Cell Localization	Cytoplasm, Cell membrane
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

WB=1:500-2000B B IHC-P=1:50-200B IHC-F=1:50-200B

Background

Protease that specifically cleaves 'Lys-63'-linked polyubiquitin chains. Has endodeubiquitinase activity. Plays an important role in the regulation of pathways leading to NF-kappa-B activation. Contributes to the regulation of cell survival, proliferation and differentiation via its effects on NF-kappa-B activation. Negative regulator of Wnt signaling. Inhibits HDAC6 and thereby promotes acetylation of alpha-tubulin and stabilization of microtubules. Plays a role in the regulation of microtubule dynamics, and thereby contributes to the regulation of cell proliferation, cell polarization, cell migration, and angiogenesis. Required for normal cell cycle progress and normal cytokinesis. Inhibits nuclear translocation of NF-kappa-B. Plays a role in the regulation of inflammation and the innate immune response, via its effects on NF-kappa-B activation. Dispensable for the maturation of intrathymic natural killer cells, but required for the continued survival of immature natural killer cells. Negatively regulates TNFRSF11A signaling and osteoclastogenesis (By similarity).

Note: This product is for in vitro research use only