## YWHAG Antibody

Catalog No: #31246

Package Size: #31246-1 50ul #31246-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_				
$\Box$	00	ori	nti	On
יע	こう	UH	บแ	on

Product Name	YWHAG Antibody		
Host Species	Rabbit		
Clonality	Polyclonal		
Applications	ELISA WB IHC		
Species Reactivity	Hu Ms Rt		
Specificity	The antibody detects endogenous level of total YWHAG protein.		
Immunogen Type	Peptide		
Immunogen Description	Synthetic peptide corresponding to a region derived from 141-154 amino acids of Human tyrosine		
	3-monooxygenase/tryptophan 5-monooxygenase activation protein, gamma polypeptide		
Target Name	YWHAG		
Other Names	tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, gamma polypeptide,		
	14-3-3GAMMA		
Accession No.	Swiss-Prot:P61981Gene ID:7532;		
Uniprot	P61981		
GeneID	7532;		
Formulation	Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.		
Storage	Store at -20°C/1 year		

## **Application Details**


Predicted MW: 28kd

ELISA: 1:2000-1:10000

Western blotting: 1:1000-1:5000

Immunohistochemistry: 1:50-1:200

## **Images**




Gel: 10%SDS-PAGE Lane1: Hela cell lysate Lane2: 293T cell lysate Lane3: Jurkat cell lysate Lysates: 40 ug per lane Primary antibody: 1/1800 dilution

Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at

1/10000 dilution

Exposure time: 10 seconds



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using 31246 (YWHAG Antibody) at dilution 1/100, on the right is treated with the synthetic peptide.

## Background

This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 100% identical to the rat ortholog. It is induced by growth factors in human vascular smooth muscle cells, and is also highly expressed in skeletal and heart muscles, suggesting an important role for this protein in muscle tissue. It has been shown to interact with RAF1 and protein kinase C, proteins involved in various signal transduction pathways.

Note: This product is for in vitro research use only