

RAD21 Antibody FITC Conjugated

Catalog No: #C05271F

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	RAD21 Antibody FITC Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt IF
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 570-620 631 derived from human Rad2
Conjugates	FITC
Target Name	RAD21
Other Names	HR21; MCD1; NXP1; SCC1; CDLS4; hHR21; HRAD21; Double-strand-break repair protein rad21 homolog; Nuclear matrix protein 1; NXP-1; SCC1 homolog; RAD21; KIAA0078
Accession No.	Swiss-Prot#O60216NCBI Gene ID5885
Uniprot	O60216
GeneID	5885;
Excitation Emission	494nm 518nm
Cell Localization	Nucleus
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=1:50-200 IF=1:50-200

Background

Cleavable component of the cohesin complex, involved in chromosome cohesion during cell cycle, in DNA repair, and in apoptosis. The cohesin complex is required for the cohesion of sister chromatids after DNA replication. The cohesin complex apparently forms a large proteinaceous ring within which sister chromatids can be trapped. At metaphase-anaphase transition, this protein is cleaved by separase ESPL1 and dissociates from chromatin, allowing sister chromatids to segregate. The cohesin complex may also play a role in spindle pole assembly during mitosis. Also plays a role in apoptosis, via its cleavage by caspase-3 CASP3 or caspase-7 CASP7 during early steps of apoptosis: the C-terminal 64 kDa cleavage product may act as a nuclear signal to initiate cytoplasmic events involved in the apoptotic pathway.

Note: This product is for in vitro research use only