

NPM (Phospho-Thr199) Antibody

Catalog No: #12031



Package Size: #12031-1 50ul #12031-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description


Product Name	NPM (Phospho-Thr199) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu Ms
Specificity	The antibody detects endogenous level of NPM only when phosphorylated at Threonine 199.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of Threonine 199 (R-D-T(p)-P-A) derived from Human NPM.
Target Name	NPM
Modification	Phospho
Other Names	B23, NPM
Accession No.	Swiss-Prot#: P06748; NCBI Gene#: 4869; NCBI Protein#: NP_001032827.1
Uniprot	P06748
GeneID	4869;
SDS-PAGE MW	32, 38kd
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C/1 year

Application Details

Predicted MW: 32B&B~38kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from Mouse heart tissue using NPM (Phospho-Thr199) Antibody #12031. The lane on the left is treated with the antigen-specific peptide.

Background

Involved in diverse cellular processes such as ribosome biogenesis, centrosome duplication, protein chaperoning, histone assembly, cell proliferation, and regulation of tumor suppressors p53/TP53 and ARF. Binds ribosome presumably to drive ribosome nuclear export. Associated with nucleolar ribonucleoprotein structures and bind single-stranded nucleic acids. Acts as a chaperonin for the core histones H3, H2B and H4. Stimulates APEX1 endonuclease activity on apurinic/aprimidinic (AP) double-stranded DNA but inhibits APEX1 endonuclease activity on AP single-stranded RNA. May exert a control of APEX1 endonuclease activity within nucleoli devoted to repair AP on rDNA and the removal of oxidized rRNA molecules. In concert with BRCA2, regulates centrosome duplication. Regulates centriole duplication: phosphorylation by PLK2 is able to trigger centriole replication. Negatively regulates the activation of EIF2AK2/PKR and suppresses apoptosis through inhibition of EIF2AK2/PKR autophosphorylation.

Note: This product is for in vitro research use only