p47 phox (Phospho-Ser359) Antibody

Catalog No: #12054

Package Size: #12054-1 50ul #12054-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| Description | |
|-----------------------|--|
| Product Name | p47 phox (Phospho-Ser359) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of p47 phox only when phosphorylated at Serine 359. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of Serine 359 |
| | (Q-R-S(p)-K-P) derived from Human p47 phox. |
| Target Name | p47 phox |
| Modification | Phospho |
| Other Names | NCF1A, NOXO2, p47phox, SH3PXD1A |
| Accession No. | Swiss-Prot#: P14598; NCBI Gene#: 653361; NCBI Protein#: NP_000256.4 |
| Uniprot | P14598 |
| GeneID | 653361; |
| SDS-PAGE MW | 45kd |
| Concentration | 1.0mg/ml |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage | Store at -20°C/1 year |

Application Details


Predicted MW: 45kd

Western blotting: 1:500~1:1000


Images



Western Blot analysis of HeLa nocodazole 1ug/ml 18h cells using Phospho-p47-phox (S359) Polyclonal Antibody diluted at 1:500



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using p47 phox (Phospho-Ser359) Antibody. The picture on the right is blocked with the phospho peptide.



Immunofluorescence analysis of HeLa cells, using p47 phox (Phospho-Ser359) Antibody. The picture on the right is blocked with the phospho peptide.

Background

NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide production).

Note: This product is for in vitro research use only