GR (Phospho-Ser203) Antibody

Catalog No: #12062

Description

Species Reactivity

Package Size: #12062-1 50ul #12062-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Product Name | GR (Phospho-Ser203) Antibody | |
|--------------|--|--|
| Host Species | Rabbit | |
| Clonality | Polyclonal | |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. | |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho | |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. | |
| Applications | WB | |
| | | |

| Immunogen Type Peptide-KLH | Specificity | The antibody detects endogenous level of GR only when phosphorylated at Serine 203. |
|----------------------------|----------------|---|
| | Immunogen Type | Peptide-KLH |

Immunogen Description Peptide sequence around phosphorylation site of Serine 203 (S-G-S(p)-P-G) derived from Human GR.

Hu

Target Name GR
Modification Phospho

Other Names GCCR, GCR, GR, GRL

Accession No. Swiss-Prot#: P04150; NCBI Gene#: 2908; NCBI Protein#: NP_000167.1

 Uniprot
 P04150

 GeneID
 2908;

 SDS-PAGE MW
 86kd

Concentration 1.0mg/ml

Formulation Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol.


Storage Store at -20°C/1 year

Application Details

Predicted MW: 86kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from A549 cells using GR (Phospho-Ser203) Antibody #12062. The lane on the left is treated with the antigen-specific peptide.

Background

Receptor for glucocorticoids (GC). Has a dual mode of action: as a transcription factor that binds to glucocorticoid response elements (GRE), both for nuclear and mitochondrial DNA, and as a modulator of other transcription factors. Affects inflammatory responses, cellular proliferation and differentiation in target tissues. Could act as a coactivator for STAT5-dependent transcription upon growth hormone (GH) stimulation and could reveal an essential role of hepatic GR in the control of body growth. Involved in chromatin remodeling. Plays a significant role in transactivation.

Note: This product is for in vitro research use only