PAK1/PAK2/PAK3(Ab-423/402/421) Antibody

Catalog No: #21169

SAB Signalway Antibody

Package Size: #21169-1 50ul #21169-2 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$		
	Accri	ıntınn
\boldsymbol{L}	๛๛	iption
		10.00


Product Name	PAK1/PAK2/PAK3(Ab-423/402/421) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB IHC IF	
Species Reactivity	Hu Ms Rt	
Specificity	The antibody detects endogenous level of total PAK1/PAK2/PAK3 protein.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around aa.421~425/400~404/419~423 (R-S-T-M-V) derived from Human	
	PAK1/PAK2/PAK3.	
Target Name	PAK1/PAK2/PAK3	
Other Names	p21-activated kinase 1; PAK-1; p65-PAK; Alpha-PAK;	
Accession No.	Swiss-Prot:Q13153Gene ID:5058	
Uniprot	Q13153	
GeneID	5058;	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

Application Details


Predicted MW: 61-67 (PAK2) 68-74 (PAK1/3)kd

Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100
Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from MCF-7 and Hela cells using PAK1/PAK2/PAK3(Ab-423/402/421) Antibody #21169.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PAK1/PAK2/PAK3(Ab-423/402/421) Antibody #21169(left) or



Immunofluorescence staining of methanol-fixed Hela cells using PAK1/PAK2/PAK3(Ab-423/402/421) Antibody #21169.

Background

The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2.

Parsons M, et al. (2005) Mol Cell Biol; 25:1680-1695. Jung JH, et al. (2005) J Biol Chem; 280:40025-40031. Lampugnani MG, et al. (2002) Mol Biol Cell; 13: 1175-1189 Wang RA, et al. (2002) EMBO J; 21: 5437 - 5447.

Note: This product is for in vitro research use only