FAK (Phospho-Ser843) Antibody

Catalog No: #11920

Package Size: #11920-1 50ul #11920-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| $\overline{}$ | | 4.5 |
|------------------|-------|-------|
| | escri | ntion |
| \boldsymbol{L} | COUL | Puon |
| | | |

| Decempation | |
|-----------------------|-----------------------------------------------------------------------------------------------------------------|
| Product Name | FAK (Phospho-Ser843) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB |
| Species Reactivity | Hu Rt |
| Specificity | The antibody detects endogenous level of FAK only when phosphorylated at serine 843. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | $Peptide\ sequence\ around\ phosphorylation\ site\ of\ serine 843 (R-G-S(p)-I-D)\ derived\ from\ Human\ FAK\ .$ |
| Target Name | FAK |
| Modification | Phospho |
| Other Names | FADK 1; FAK1; Focal adhesion kinase 1; PTK2; pp125FAK |
| Accession No. | Swiss-Prot#: Q05397; NCBI Gene#: 5747; NCBI Protein#: NP_001186578.1 |
| Uniprot | Q05397 |
| GeneID | 5747; |
| SDS-PAGE MW | 120kd |
| Concentration | 1.0mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide |
| | and 50% glycerol. |
| Storage | Store at -20°C/1 year |
| | |

Application Details

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from 3T3 cells treated with PMA using Phospho-FAK (Ser843) antibody #11920.The lane on the right is treated with the antigen-specific peptide.

Background

Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity.

Jiang X, Sinnett-Smith J, Rozengurt E (2007)Cell Signal 19, 1000-10 . Jacamo R, Jiang X, Lunn JA, Rozengurt E (2007)J Cell Physiol 210, 436-44. Le Boeuf F, Houle F, Sussman M, Huot J (2006)Mol Biol Cell 17, 3508-20 .

Note: This product is for in vitro research use only