

## IKK- alpha/ beta (Phospho-Ser176/177) Antibody

Catalog No: #11931



Package Size: #11931-1 50ul #11931-2 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)


## Description

Product Name	IKK- alpha/ beta (Phospho-Ser176/177) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB,IHC,IF,ELISA
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of IKK- alpha/beta only when phosphorylated at serine 176/177.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 176/177 (Q-G-S(p)-L-C) derived from Human IKK-alpha/beta.
Target Name	IKK- alpha/ beta
Modification	Phospho
Other Names	FLJ40509; I-kappa-B kinase; IKBKB; kinase beta; NFKB1KB
Accession No.	Swiss-Prot#: O15111/O14920; NCBI Gene#: 1147; NCBI Protein#: NP_001269.3
Uniprot	O15111
GeneID	1147;
SDS-PAGE MW	85kd
Concentration	1.0mg/ml
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C/1 year

## Application Details

Western blotting: 1:500~1:1000

## Images



Western blot analysis of extracts from NIH-3T3 cells treated with TNF using IKK- alpha/ beta (Phospho-Ser176/177) antibody #11931. The lane on the right is treated with the antigen-specific peptide.

## Background

Acts as part of the IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. Also phosphorylates NCOA3.

Chandrakesan P, et al. (2010) J Biol Chem 285, 33485-98

Hinz M, et al. (2010) Mol Cell 40, 63-74

Choudhary S, Lu M, Cui R, Brasier AR (2007) Mol Endocrinol 21, 2203-17

Note: This product is for in vitro research use only