## ETS1 (Phospho-Thr38) Antibody

Catalog No: #11658

Package Size: #11658-1 50ul #11658-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


	മല	۱'n	$\sim$	$\mathbf{a}$	n
$\boldsymbol{L}$	esc	/I II	υu	v	
		١ ١			

Product Name	ETS1 (Phospho-Thr38) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.	
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho	
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.	
Applications	WB IHC IF	
Species Reactivity	Hu Ms	
Specificity	The antibody detects endogenous levels of ETS1 only when phosphorylated at threonine 38.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around phosphorylation site of threonine 38(L-L-T(p)-P-S) derived from Human ETS1 .	
Target Name	ETS1	
Modification	Phospho	
Other Names	p54; ETS1; ETS-1; C-ets-1 protein;	
Accession No.	Swiss-Prot#: P27577; NCBI Gene#: 2113; NCBI Protein#: NP_035938.2.	
Uniprot	P27577	
GeneID	23871;	
SDS-PAGE MW	54kd	
Concentration	1.0mg/ml	
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide	
	and 50% glycerol.	
Storage	Store at -20°C/1 year	


## Application Details

Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100
Immunofluorescence: 1:100~1:200


## **Images**



Western blot analysis of extracts from HepG2 cells using ETS1 (Phospho-Thr38) Antibody #11658. The lane on the right is treated with the antigen-specific peptide.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue, using ETS1 (Phospho-Thr38) antibody #11658 (left)or the same antibody preincubated with blocking peptide (right).



Immunofluorescence staining of methanol-fixed HeLa cells using ETS1 (Phospho-Thr38) Antibody #11658.

## Background

Transcription factor. Directly controls the expression of cytokine and chemokine genes in a wide variety of different cellular contexts. May control the differentiation, survival and proliferation of lymphoid cells. May also regulate angiogenesis through regulation of expression of genes controlling endothelial cell migration and invasion.

Tsukasa Higuchi, Mol. Cell. Biol., May 2007; 27: 3353 - 3366.

NK Bhat, PNAS, May 1990; 87: 3723.

H Suzuki, PNAS, May 1995; 92: 4442.

V Fafeur, Cell Growth Differ., Jun 1997; 8: 655.

Note: This product is for in vitro research use only