## **ICOS** Antibody

Catalog No: #32445

SAB Signalway Antibody

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Package Size: #32445-1 50ul #32445-2 100ul

Host Species Rabbit  Clonality Polyclor  Purification Antibod  Applications WB,IF  Species Reactivity Human,  Specificity The ant	
Clonality Polyclor Purification Antibod Applications WB,IF Species Reactivity Human, Specificity The ant	dies were purified by affinity purification using immunogen.  a,Mouse,Rat tibody detects endogenous level of total ICOS protein.
Purification Antibod  Applications WB,IF  Species Reactivity Human,  Specificity The ant	dies were purified by affinity purification using immunogen.  a,Mouse,Rat tibody detects endogenous level of total ICOS protein.
Applications WB,IF Species Reactivity Human, Specificity The ant	i,Mouse,Rat tibody detects endogenous level of total ICOS protein.
Species Reactivity Human, Specificity The ant	tibody detects endogenous level of total ICOS protein.
Specificity The ant	tibody detects endogenous level of total ICOS protein.
· · · ·	·
	binant Protein
Immunogen Type Recomb	
Immunogen Description Recomb	binant protein of human ICOS.
Target Name ICOS	
Other Names AILIM; 0	CD278; CVID1; MGC39850;
Accession No. Swiss-F	Prot:Q9Y6W8NCBI Gene ID:29851
Uniprot Q9Y6W	V8
GeneID 29851;	
SDS-PAGE MW 23KD	
Concentration 1.0mg/r	ml
Formulation Supplie	ed at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
sodium	azide and 50% glycerol.
Storage Store at	at -20°C


## Application Details

WB□1:500 - 1:2000IF□1:50 - 1:200


## Images




Immunofluorescence analysis of rat thymus using ICOS at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of mouse thymus using ICOS at dilution of 1:100. Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using ICOS at 1:1000 dilution.



Immunofluorescence analysis of human spleen using ICOS at dilution of 1:100. Blue: DAPI for nuclear staining.

## Background

The protein encoded by this gene belongs to the CD28 and CTLA-4 cell-surface receptor family. It forms homodimers and plays an important role in cell-cell signaling, immune responses, and regulation of cell proliferation.

Note: This product is for in vitro research use only