## p56Dok-2(Ab-299) Antibody

Catalog No: #21270

Package Size: #21270-1 50ul #21270-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| _ |       | 4.5   |    |
|---|-------|-------|----|
|   | escri | Intra | าท |
|   |       | IVIII |    |
|   |       |       | ٠  |


| The state of the s |   |
|--|---|
| Product Name   | p56Dok-2(Ab-299) Antibody   |
| Host Species   | Rabbit  |
| Clonality  | Polyclonal  |
| Purification   | Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were |
|  | purified by affinity-chromatography using epitope-specific peptide.                                       |
| Applications   | WB IHC IF   |
| Species Reactivity   | Hu  |
| Specificity  | The antibody detects endogenous level of total p56Dok-2 protein.  |
| Immunogen Type   | Peptide-KLH   |
| Immunogen Description  | Peptide sequence around aa.297~301 (G-E-Y-A-V) derived from Human p56Dok-2.                               |
| Target Name  | p56Dok-2  |
| Other Names  | DOK2  |
| Accession No.  | Swiss-Prot: O60496NCBI Protein: NP_003965.2   |
| Uniprot  | O60496  |
| GeneID   | 9046;   |
| Concentration  | 1.0mg/ml  |
| Formulation  | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%      |
|  | sodium azide and 50% glycerol.  |
| Storage  | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.                 |
|  |   |

## Application Details


Predicted MW: 56kd

Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100
Immunofluorescence: 1:100~1:200


## **Images**



Western blot analysis of extracts from JK cells using p56Dok-2(Ab-299) Antibody #21270.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using p56Dok-2(Ab-299) Antibody #21270(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using p56Dok-2(Ab-299) Antibody #21270.

## Background

DOK proteins are enzymatically inert adaptor or scaffolding proteins. They provide a docking platform for the assembly of multimolecular signaling complexes. DOK2 may modulate the cellular proliferation induced by IL-4, as well as IL-2 and IL-3. May be involved in modulating Bcr-Abl signaling. Attenuates EGF-stimulated MAP kinase activation

Feng Cong, et,al. (1999) Mol. Cell. Biol; 19: 8314 - 8325.

Serge Lemay, et,al. (2000) Mol. Cell. Biol; 20: 2743 - 2754.

Ute Schaeper, et,al.(2000) J. Cell Biol; 149: 1419.

Miyuki Honma, et,al. (2006) Genes Cells; 11: 143 - 151.

Note: This product is for in vitro research use only