## syk(Ab-323) Antibody

Catalog No: #21546

Package Size: #21546-1 50ul #21546-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_		4.5	
Desc	rın	tin	n
レしるし	пρ	เมษ	ш


Product Name	syk(Ab-323) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB IHC IF	
Species Reactivity	Hu	
Specificity	The antibody detects endogenous level of total syk protein.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around aa. 321~325 (N-P-Y-E-P) derived from Human syk.	
Target Name	syk	
Other Names	Spleen tyrosine kinase	
Accession No.	Swiss-Prot: P43405NCBI Protein: NP_001128524.1	
Uniprot	P43405	
GeneID	6850;	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

## Application Details


Predicted MW: 72kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


## **Images**



Western blot analysis of extracts from JK cells using syk(Ab-323) Antibody #21546.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using syk(Ab-323) Antibody #21546(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using syk(Ab-323) Antibody #21546.

## Background

Positive effector of BCR-stimulated responses. Couples the B-cell antigen receptor (BCR) to the mobilization of calcium ion either through a phosphoinositide 3-kinase-dependent pathway, when not phosphorylated on tyrosines of the linker region, or through a phospholipase C-gamma-dependent pathway, when phosphorylated on Tyr-348 and Tyr-352. Thus the differential phosphorylation of Syk can determine the pathway by which BCR is coupled to the regulation of intracellular calcium ion

Zhang, J. et al. (2000) J. Biol. Chem. 275, 35442-35447.

Turner, M. et al. (2000) Immunol. Today 21, 148-154.

Decker, M. et al. (1998) J. Biol. Chem. 273, 8867-8874.

Law, C.L. et al. (1996) Mol. Cell. Biol. 16, 1305-1315.

Note: This product is for in vitro research use only