

CD44 Polyclonal Antibody Cy7 Conjugated

Catalog No: #C04151Cy7

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	CD44 Polyclonal Antibody Cy7 Conjugated
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Purified by Protein A.
Applications	Flow-Cyt
Species Reactivity	Hu Ms Rt
Immunogen Description	KLH conjugated synthetic peptide aa 671-710 742 derived from human CD44
Conjugates	Cy7
Target Name	CD44
Other Names	CD44 antigen; CDw44; Epican; Extracellular matrix receptor III; ECMR-III; GP90 lymphocyte homing adhesion receptor; HUTCH-I; Heparan sulfate proteoglycan; Hermes antigen; Hyaluronate receptor; Phagocytic glycoprotein 1; PGP-1; Phagocytic glycoprotein I; PGP-I; CD44; LHR; MDU2; MDU3; MIC4
Accession No.	Swiss-Prot#:P16070NCBI Gene ID:960
Uniprot	P16070
GeneID	960;
Excitation Emission	743nm 767nm
Cell Localization	Cytoplasm
Concentration	1mg ml
Formulation	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Application Details

Flow-Cyt=2ug/Test

Background

Receptor for hyaluronic acid (HA). Mediates cell-cell and cell-matrix interactions through its affinity for HA, and possibly also through its affinity for other ligands such as osteopontin, collagens, and matrix metalloproteinases (MMPs). Adhesion with HA plays an important role in cell migration, tumor growth and progression. In cancer cells, may play an important role in invadopodia formation. Also involved in lymphocyte activation, recirculation and homing, and in hematopoiesis. Altered expression or dysfunction causes numerous pathogenic phenotypes. Great protein heterogeneity due to numerous alternative splicing and post-translational modification events.

Note: This product is for in vitro research use only