fumarate hydratase antibody

Catalog No: #22113



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$			400	
	esc	rın	tic	ı'n
\boldsymbol{L}	しつし	IIIU	uu	и I

Product Name	fumarate hydratase antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC IF
Species Reactivity	Hu
Immunogen Type	Recombinant protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 195 and 443
	of fumarate hydratase
Target Name	fumarate hydratase
Accession No.	Swiss-Prot:P07954Gene ID:2271
Uniprot	P07954
GeneID	2271;
Concentration	0.9mg/ml
Formulation	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a
	preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 55kd

Western blotting: 1:500-1:3000

Immunohistochemistry: 1:100-1:500

Immunofluorescence: 1:100-1:200

Images




Sample (30 ug of whole cell lysate)


A: HeLa B: Hep G2

10% SDS PAGE

Primary antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded NCIN87 xenograft, using fumarate hydratase antibody at 1: 500 dilution.



Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using fumarate hydratase antibody at 1: 200 dilution.

Background

The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy. [provided by RefSeq]

Note: This product is for in vitro research use only