## ZHX2 antibody

Catalog No: #22147



Orders: order@signalwayantibody.com Support: tech@signal way antibody.com

$\overline{}$			
	escr	מוי	tion
$\boldsymbol{L}$	COUL	ıv	เเบเ

ZHX2 antibody	
Rabbit	
Polyclonal	
Purified by antigen-affinity chromatography.	
WB IHC IF	
Hu	
Recombinant protein	
Recombinant protein fragment contain a sequence corresponding to a region within amino acids 533 and 796	
of ZHX2	
ZHX2	
Swiss-Prot:Q9Y6X8Gene ID:22882	
Q9Y6X8	
22882;	
0.9mg/ml	
Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a	
preservative.	
Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

## Application Details


Predicted MW: 92kd

Western blotting: 1:500-1:3000

Immunohistochemistry: 1:100-1:250


Immunofluorescence: 1:100-1:200

## **Images**




Sample (30 ug of whole cell lysate) A: Hep G2 7.5% SDS PAGE

Primary antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded SAS Xenograft, using ZHX2 antibody at 1: 100 dilution.



Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using ZHX2 antibody at 1: 200 dilution.

## Background

The members of the zinc fingers and homeoboxes gene family are nuclear homodimeric transcriptional repressors that interact with the A subunit of nuclear factor-Y (NF-YA) and contain two C2H2-type zinc fingers and five homeobox DNA-binding domains. This gene encodes member 2 of this gene family. In addition to forming homodimers, this protein heterodimerizes with member 1 of the zinc fingers and homeoboxes family. [provided by RefSeq]

Note: This product is for in vitro research use only