Product Datasheet

MEK2(Phospho-Thr394) Antibody

Catalog No: #11008

Package Size: #11008-1 50ul #11008-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| | escri | ال ا م | | ı |
|------------------|-------|---------------|--------------|---|
| | escri | DТ | or | ۱ |
| $\boldsymbol{-}$ | COUL | ν u | \mathbf{o} | Ц |


| Product Name | MEK2(Phospho-Thr394) Antibody | |
|-----------------------|--------------------------------------------------------------------------------------------------------|--|
| Host Species | Rabbit | |
| Clonality | Polyclonal | |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. | |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho | |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. | |
| Applications | WB IHC IF | |
| Species Reactivity | Hu | |
| Specificity | The antibody detects endogenous level of MEK2 only when phosphorylated at threonine 394. | |
| Immunogen Type | Peptide-KLH | |
| Immunogen Description | Peptide sequence around phosphorylation site of threonine 394 (P-G-T(p)-P-T) derived from Human MEK-2. | |
| Conjugates | Unconjugated | |
| Target Name | MEK2 | |
| Modification | Phospho | |
| Other Names | ERK activator kinase 2; MAP kinase kinase 2; MAP2K2; MAPK/ERK kinase 2; MAPKK 2 | |
| Accession No. | Swiss-Prot: P36507NCBI Protein: NP_109587.1 | |
| Concentration | 1.0mg/ml | |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% | |
| | sodium azide and 50% glycerol. | |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. | |
| | | |

Application Details


Predicted MW: 44kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from HepG2 and Hela cells untreated or treated with UV using MEK-2(Phospho-Thr394) Antibody #11008.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using MEK-2(Phospho-Thr394) Antibody #11008(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using MEK-2(Phospho-Thr394) Antibody #11008.

Background

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases.

Crews C M, et al. (1992) Science. 258:478-480.

Alessi D R, et al. (1994) EMBO J. 13:1610-1619.

Rosen L B, et al. (1994) Neuron. 12:1207-1221.

Cowley S, et al. (1994) Cell. 77:841-852.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.