PKD/PKCm(Phospho-Ser910) Antibody

Catalog No: #11096

Package Size: #11096-1 50ul #11096-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| $\overline{}$ | | |
|------------------|-------|----------|
| | Accri | ption |
| \boldsymbol{L} | ๛๛ | ווטווטוו |
| | | 10.00 |

| Product Name | PKD/PKCm(Phospho-Ser910) Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB IF |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of PKD/PKCm only when phosphorylated at serine 910. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 910 (R-V-S(p)-I-L) derived from Human PKD/PKCm. |
| Conjugates | Unconjugated |
| Target Name | PKD/PKCm |
| Modification | Phospho |
| Other Names | KPCD1; PKC-mu; PKCM; PKD; PRKCM |
| Accession No. | Swiss-Prot: Q15139NCBI Protein: NP_002733.2 |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |


Application Details

Predicted MW: 115kd


Western blotting: 1:500~1:1000

Immunofluorescence: 1:100~1:200

Images



Western blot analysis of extracts from EGF-treated 293 and PMA-treated HUVEC cells using PKD/PKCm(Phospho-Ser910) Antibody #11096.



Immunofluorescence staining of methanol-fixed Hela cells using PKD/PKCm(Phospho-Ser910) Antibody #11096.

Background

Converts transient diacylglycerol. (DAG) signals into prolonged physiological effects, downstream of PKC. Involved in resistance to oxidative stress through activation of NF-kappa-B.

Matthews SA, et al. J Biol Chem 1999 Sep; 274(37): 26543-26549

Brandlin I, et al. J Biol Chem 2002 Feb; 277(8): 6490-6496 Storz P, et al. Mol Pharmacol 2004 Oct; 66(4): 870-879

Note: This product is for in vitro research use only and is not intended for use in humans or animals.