EGFR(Phospho-Tyr869) Antibody

Catalog No: #11229

Package Size: #11229-1 50ul #11229-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


| Product Name | EGFR(Phospho-Tyr869) Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous level of EGFR only when phosphorylated at tyrosine 869. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of tyrosine 869 (K-E-Y(p)-H-A) derived from Human EGFR. |
| Conjugates | Unconjugated |
| Target Name | EGFR |
| Modification | Phospho |
| Other Names | ERBB1; Receptor protein-tyrosine kinase ErbB-1; kinase EGFR |
| Accession No. | Swiss-Prot: P00533NCBI Protein: NP_005219.2 |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |

Application Details

Predicted MW: 175kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from MDA cells untreated or treated with EGF using EGFR(Phospho-Tyr869) Antibody #11229.

Background

Receptor for EGF, but also for other members of the EGF family, as TGF-a, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. Phosphorylates MUC1 in breast cancer cells and increases the interaction of MUC1 with SRC and CTNNB1/beta-catenin.

Inoue A, et al. (2005) PLoS Med; 2(1): e13 Sun H, et al. (2004) EMBO J; 23(1): 100-110

Kanner SB, et al. (1991) Mol Cell Biol; 11(2): 713-720 Wu TT, et al. (1998) Mol Biol Cell; 9(7): 1661-1674

0

Published Papers

el at., EGFR-Phosphorylated Platelet Isoform of Phosphofructokinase 1 Promotes PI3K Activation. IN Mol Cell.On 2018 Apr 19 by Lee JH, Liu R et al..PMID:29677490, , (2018)

PMID:29677490

el at., Secreted and O-GlcNAcylated MIF binds to the human EGF receptor and inhibits its activation. In Nat Cell Biol on 2015 Oct by Yanhua Zheng, Xinjian Li et al.. PMID:26280537 , , (2015)

PMID:26280537

el at., Mutual regulation between phosphofructokinase 1 platelet isoform and VEGF promotes glioblastoma tumor growth. In Cell Death Dis on 2022 Nov 26 by Je Sun Lim, YuJie Shi, et al..PMID:36435833, , (2022)

PMID:36435833

Note: This product is for in vitro research use only and is not intended for use in humans or animals.