# STAT1(Ab-701) Antibody

Catalog No: #21044

Package Size: #21044-1 50ul #21044-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

# Description


Product Name	STAT1(Ab-701) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were
	purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total STAT1 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa. 699~703 (T-G-Y-I-K) derived from Human STAT1.
Conjugates	Unconjugated
Target Name	STAT1
Other Names	CANDF7, ISGF-3, STAT91
Accession No.	Swiss-Prot: P42224NCBI Protein: NP _009330.1
SDS-PAGE MW	84,91kd
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

# **Application Details**


Predicted MW: 84,91kd Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

# **Images**



Western blot analysis of extracts from Hela, A431 and JK cells using STAT1(Ab-701) Antibody #21044.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using STAT1(Ab-701) Antibody #21044(left) or the same antibody preincubated with blocking peptide(right).

# Background

Signal transducer and activator of transcription that mediates signaling by interferons (IFNs). Following type I IFN (IFN-a and IFN-beta) binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state. In response to type II IFN (IFN-gamma), STAT1 is tyrosine- and serine-phosphorylated. It then forms a homodimer termed IFN-gamma-activated factor (GAF), migrates into the nucleus and binds to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state.

Heim M H, (1999) J Recept Signal Transduct Res. 19: 75-120.

Durbin J E, et al. (1996) Cell. 84: 443-450.

Meraz M A, et al. (1996) Cell. 84: 431-442.

Wakao H, et al. (1994) EMBO J. 13: 2182-2191.

Demoulin J, B. et al. (1999) J Biol Chem. 274: 25855-258

Ihle J N, et al. (1994) Trends Biochem Sci. 19: 222-227.

### **Published Papers**

Xiang Cheng, Jing Wang, Ni Xia el at., A guanidine-rich regulatory oligodeoxy-nucleotideimproves type-2 diabetes in obese mice by blocking T-cell differentiation., EMBO Mol Med., 4(10):1112η— C1125(2012)

#### PMID:23027613

el at., Danzhi Jiangtang Capsule ameliorates kidney injury via inhibition of the JAK-STAT signaling pathway and increased antioxidant capacity in STZ-induced diabetic nephropathy rats.In Biosci Trends.On 2019 Jan 22 by Sun M, Bu W et al..PMID: 30606979, (2018)

#### PMID:30606979

el at., A thiazole-derived oridonIn analogue exhibits antitumor activity by directly and allosterically InhibitIng STAT3. In J Biol Chem on 2019 Nov 15;by Shen X, Zhao L,et al..PMID:31594861, , (2019)

#### PMID:31594861

el at., A Guanidine-Rich Regulatory Oligodeoxynucleotide Improves type-2 Diabetes in Obese Mice by Blocking T-cell Differentiation. In EMBO Mol Med on 2012 Oct by Xiang Cheng, Jing Wang, et al..PMID:23027613

, , (2012)

#### PMID:23027613

el at., DJC suppresses Advanced Glycation End Products-Induced JAK-STAT signaling and ROS in mesangial cells.In Evid Based Complement Alternat Med on 2017 May 29 by Min Sun, Yan Li ,et al..PMID: 28630633, , (2017)

### PMID:28630633

el at., Expression of interferon effector gene SART1 correlates with interferon treatment response against hepatitis B infection. In Mediators Inflamm on 2016 by Yong Li, Chuanlong Zhu et al..PMID: 28077916, , (2016)

#### PMID:28077916

el at., ycobacterium tuberculosis EspB protein suppresses interferon-ηΊ¬-induced autophagy in murine macrophages.In J Microbiol Immunol Infect on 2016 Dec by DanDan Huang, Lang Bao et al..PMID:25641595 , , (2016)

#### PMID:25641595

el at., IL-17 intensifies IFN-ŋ1¬-induced NOS2 upregulation in RAW 264.7 cells by further activating STAT1 and NF-I-• B.In Int J Mol Med on 2016 byQiyue Gao, Yang Liu et al..PMID:26677135 , , (2016)

#### PMID:26677135

el at., Costunolide improved dextran sulfate sodium-induced acute ulcerative colitis in mice through NF-kB, STAT1/3, and Akt signaling pathways. In Int Immunopharmacol on 2020 Jul by Fan Xie, Hai Zhang, et al..PMID:32413737, , (2020)

PMID:32413737

Note: This product is for in vitro research use only and is not intended for use in humans or animals.