

IRF1 Rabbit mAb

Catalog No: #48646

Package Size: #48646-1 50ul #48646-2 100ul


Description

Product Name	IRF1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SR44-08
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	Interferon regulatory factor 1 antibody Interferon regulatory factor 1 isoform +I9 antibody Interferon regulatory factor 1 isoform d78 antibody Interferon regulatory factor 1 isoform delta4 antibody Interferon regulatory factor 1 isoform delta7 antibody IRF 1 antibody IRF-1 antibody IRF1 antibody IRF1_HUMAN antibody MAR antibody MAR1 antibody
Accession No.	Swiss-Prot#:P10914
Calculated MW	48 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of IRF1 on different lysates using anti-IRF1 antibody at 1/1,000 dilution. Positive control:
Lane 1: PC-12
Lane 2: Jurkat




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue using anti-IRF1 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-IRF1 antibody. Counter stained with hematoxylin.



ICC staining IRF1 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining IRF1 in NIH/3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Jurkat cells with IRF1 antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding factors that function as regulators of both type I interferon (interferon- and) and interferon-inducible genes. The two factors are structurally related, particularly in their N-terminal regions, which confer DNA binding specificity. In addition, both bind to the same sequence within the promoters of interferon- and interferon- genes. IRF-1 functions as an activator of interferon transcription, while IRF-2 binds to the same cis elements and represses IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in regulating cell growth; overexpression of the repressor IRF-2 leads to cell transformation while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7, ISGF-3 p48 (a component of the ISGF-3 complex) and IFN consensus sequence-binding protein (ICSBP).

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.