

Cdc27 Rabbit mAb

Catalog No: #49210

Package Size: #49210-1 50ul #49210-2 100ul



SAB[®]
Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Cdc27 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SD85-02
Purification	ProA affinity purified
Applications	WB
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	ANAPC3 antibody Anaphase Promoting Complex 3 antibody Anaphase promoting complex protein 3 antibody Anaphase Promoting Complex Subunit 3 antibody Anaphase-promoting complex subunit 3 antibody APC 3 antibody APC3 antibody Cdc 27 antibody Cdc27 antibody CDC27 homolog antibody CDC27_HUMAN antibody CDC27Hs antibody Cell Division Cycle 27 antibody Cell division cycle protein 27 homolog antibody D0S1430E antibody D17S978E antibody H NUC antibody H-NUC antibody HNUC antibody Nuc2 homolog antibody
Accession No.	Swiss-Prot#:P30260
Calculated MW	92 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:1,000-5,000

Images



Western blot analysis of Cdc27 on different lysates using anti-Cdc27 antibody at 1/1,000 dilution. Positive control:
Lane 1: HeLa Lane 2: K562

Background

n the cell cycle. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. Cdc6 is the human homolog of *Saccharomyces cerevisiae* Cdc6, which is involved in the initiation of DNA replication. Cdc37 appears to facilitate Cdk4/cyclin D1 complex formation and has been shown to form a stable complex with Hsp90. Cdc34, Cdc27 and Cdc16 function as ubiquitin-conjugating enzymes. Cdc34 is thought to be the structural and functional homolog of *Saccharomyces cerevisiae* Cdc34, which is essential for the G1 to S phase transition. Cdc16 and Cdc27 are components of the APC (anaphase-promoting complex) which ubiquitinates cyclin B, resulting in cyclin B/Cdk complex degradation.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.