

SUZ12 Rabbit mAb

Catalog No: #49274

Package Size: #49274-1 50ul #49274-2 100ul



SAB[®]
Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	SUZ12 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JJ09-04
Purification	ProA affinity purified
Applications	WB, ICC/IF, IP
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	ChET 9 protein antibody CHET9 antibody Chromatin precipitated E2F target 9 protein antibody JJAZ1 antibody Joined to JAZF1 protein antibody KIAA0160 antibody Polycomb protein SUZ12 antibody Suppressor of zeste 12 homolog antibody Suppressor of zeste 12 protein homolog antibody SUZ12 antibody SUZ12 polycomb repressive complex 2 subunit antibody SUZ12_HUMAN antibody
Accession No.	Swiss-Prot#:Q15022
Calculated MW	83 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 ICC: 1:50-1:200


Images




Western blot analysis of SUZ12 on MCF-7 cells lysates using anti-SUZ12 antibody at 1/1,000 dilution.




ICC staining SUZ12 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining SUZ12 in MCF-7 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining SUZ12 in SW480 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining SUZ12 in SKOV-3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

The Polycomb group (PcG) genes contribute to the maintenance of cell identity, cell cycle regulation and oncogenesis. The mammalian PcG proteins are regulatory proteins important for Hox gene expression, axial skeleton development and the control of proliferation and survival of hematopoietic cells. By inducing changes in chromatin structure, the PcG proteins are part of a cellular memory system that is responsible for gene activity being inherited to progeny cells. PcG proteins silence gene expression through the formation of multimeric protein complexes with different compositions. Manipulating the expression-levels of various PcG proteins in mammalian cell lines results in cellular transformation, which may be a link between the chromatin-associated PcG proteins and cancer. Polycomb protein SUZ12, also designated ChET 9 protein or joined-to-JAZF1 protein, is a nuclear protein belonging to the VEFS (VRN2-EMF2-FIS2-SUZ12) family. SUZ12 has been detected at the breakpoints of a certain recurrent chromosomal

translocation which has been reported in endometrial stromal sarcoma. It is a component of the PRC2 complex, composed of EED, EZH2, SUZ12/JJAZ1, RBBP4 and RBBP7.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.