

CDC42 Rabbit mAb

Catalog No: #49282

Package Size: #49282-1 50ul #49282-2 100ul

Description


Product Name	CDC42 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JJ086-04
Purification	ProA affinity purified
Applications	WB IHC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	CDC42 antibody CDC42_HUMAN antibody CDC42Hs antibody Cell division control protein 42 homolog antibody Cell division cycle 42 (GTP binding protein 25kDa) antibody Cell division cycle 42 antibody dJ224A6.1.1 (cell division cycle 42 (GTP-binding protein, 25kD)) antibody dJ224A6.1.2 (cell division cycle 42 (GTP-binding protein, 25kD)) antibody G25K antibody G25K GTP-binding protein antibody Growth regulating protein antibody GTP binding protein 25kDa antibody Small GTP binding protein CDC42 antibody TKS antibody
Accession No.	Swiss-Prot#:P60953
Calculated MW	Predicted band size: 21 kDa
SDS-PAGE MW	Observed band size: 21 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:2000

IHC: 1:50-1:200

Images



All lanes: CDC42 Rabbit mAb at 1/1k dilution

Lane 1 : Hela whole cell lysates
Lane 2 : JK whole cell lysates
Lane 3 : C6 whole cell lysates
Lane 4 : Mouse brain lysates
Lane 5 : Rat brain lysates

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 21 kDa

Observed band size: 21 kDa

Exposure time: 3 seconds




All lanes: CDC42 Rabbit mAb at 1/1k dilution


Lane 1 : Wild-type HeLa cell lysate

Lane 2 : CDC42 knockdown HeLa cell lysate


Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human colon tissue stained for CDC42 using 49282 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded mouse skin tissue stained for CDC42 using 49282 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded rat skin tissue stained for CDC42 using 49282 at 1/100 dilution in immunohistochemical analysis.

Background

The superfamily of GTP-binding proteins, for which the Ras proteins are prototypes, has been implicated in regulation of diverse biological activities involving various aspects of cell growth and division. One mammalian member of the family, Cdc42, has an amino acid sequence that is similar to those of various members of the Ras superfamily proteins, including N-, K- and H-Ras, Rho proteins and the Rac proteins. On the basis of in vitro phosphorylation studies, it has been suggested that human Cdc42 may function in the signaling pathway of the EGF receptor or related growth factor receptor protein kinases. The Dbl oncogene has been shown to specifically catalyze dissociation of GDP from human Cdc42.

Published Papers

el at., A transcribed ultraconserved noncoding RNA, uc.285+, promotes colorectal cancer proliferation through dual targeting of CDC42 by directly binding mRNA and protein. In Transl Res on 2024 Aug by Sixian Chen, Qingyun Zhao,et al..PMID:38552953, , (2024)

[PMID:38552953](#)

Chen Sixian;Zhao Qingyun;Zhang Ruirui;Liu Jungang;Peng Wenyi;Xu Haotian;Li Xiaofei;Wang Xin;Wu Shuilian;Li Gang;Nan Aruo el at., A transcribed ultraconserved noncoding RNA, uc.285+, promotes colorectal cancer proliferation through dual targeting of CDC42 by directly binding mRNA and protein, , (2024)

[PMID:](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.